

Determining α_s and nPDFs from jets in DIS and photoproduction

Michael Klasen

Institute for Theoretical Physics, University of Münster

16 November 2016

Work done in collaboration with T. Biekötter, G. Kramer and M. Michael



References

Two recent publications:

- MK, G. Kramer, M. Michael
NNLO contributions to jet photoproduction and determination of α_s
Phys. Rev. D 89 (2014) 074032 [arXiv:1310.1724]
- T. Biekötter, MK, G. Kramer
NNLO contributions to inclusive jet production in DIS and
determination of α_s
Phys. Rev. D 92 (2015) 074037 [arXiv:1508.07153]

References

Referring to two final HERA publications:

- H. Abramowicz et al. [ZEUS Collaboration]
Inclusive-jet photoproduction at HERA and determination of α_s
Nucl. Phys. B 864 (2012) 1
- V. Andreev et al. [H1 Collaboration]
Measurement of multijet production in ep collisions at high Q^2
and determination of the strong coupling α_s
Eur. Phys. J. C 75 (2015) 65

References

Referring to two final HERA publications:

- H. Abramowicz et al. [ZEUS Collaboration]
Inclusive-jet photoproduction at HERA and determination of α_s
Nucl. Phys. B 864 (2012) 1
- V. Andreev et al. [H1 Collaboration]
Measurement of multijet production in ep collisions at high Q^2
and determination of the strong coupling α_s
Eur. Phys. J. C 75 (2015) 65

Upcoming:

- V. Andreev et al. [H1 Collaboration]
Measurement of jet production cross sections in deep-inelastic ep
scattering at HERA
DESY 16-200, to be submitted to Eur. Phys. J. C

Unified approach to NNLO soft and virtual corrections

N. Kidonakis, Int. J. Mod. Phys. A 19 (2004) 1793

- Full NNLO calculations challenging, slowly making progress
- Soft/virtual corrections often dominant, e.g. close to threshold

$$z \equiv \frac{(p_1 + p_2)^2}{(p_a + p_b)^2} \rightarrow 1$$

- Resummation of these corrections possible to all orders
- Reexpansion gives approximate NNLO (aNNLO) results
- Results depend on 1PI or PIM kinematics, $\overline{\text{MS}}$ or DIS scheme

NLO master formula

$$\begin{aligned}
 d\sigma_{ab} &= d\sigma_{ab}^B \frac{\alpha_s(\mu)}{\pi} [c_3 D_1(z) + c_2 D_0(z) + c_1 \delta(1-z)] \\
 &+ \frac{\alpha_s^{d_{\alpha_s}+1}(\mu)}{\pi} [A^c D_0(z) + T_1^c \delta(1-z)]
 \end{aligned}$$

$$D_l(z) = \left[\frac{\ln^l(1-z)}{1-z} \right]_+$$

$$d_{\alpha_s} = 0, 1, 2, \dots, \text{ if Born is of } \mathcal{O}(\alpha_s^{0,1,2,\dots})$$

Leading coefficients (simple color flow)

QCD Compton process: $\gamma q \rightarrow qg$

$$c_3 = C_F - N_C,$$

$$c_2 = C_F \left[-\ln \left(\frac{\mu_p^2}{s} \right) - \frac{3}{4} + 2 \ln \left(\frac{-u}{s} \right) \right] + N_C \ln \left(\frac{t}{u} \right) - \frac{\beta_0}{4},$$

$$c_1^\mu = -\frac{3C_F}{4} \ln \left(\frac{\mu_p^2}{s} \right) + \frac{\beta_0}{4} \ln \left(\frac{\mu^2}{s} \right)$$

Leading coefficients (simple color flow)

QCD Compton process: $\gamma q \rightarrow qg$

$$c_3 = C_F - N_C,$$

$$c_2 = C_F \left[-\ln \left(\frac{\mu_p^2}{s} \right) - \frac{3}{4} + 2 \ln \left(\frac{-u}{s} \right) \right] + N_C \ln \left(\frac{t}{u} \right) - \frac{\beta_0}{4},$$

$$c_1^\mu = -\frac{3C_F}{4} \ln \left(\frac{\mu_p^2}{s} \right) + \frac{\beta_0}{4} \ln \left(\frac{\mu^2}{s} \right)$$

Photon gluon fusion: $\gamma g \rightarrow q\bar{q}$

$$c_3 = 2(N_C - C_F),$$

$$c_2 = -\frac{3C_F}{2} + N_C \left[-\ln \left(\frac{\mu_p^2}{s} \right) + \ln \left(\frac{tu}{s^2} \right) \right],$$

$$c_1^\mu = -\frac{\beta_0}{4} \ln \left(\frac{\mu_p^2}{s} \right) + \frac{\beta_0}{4} \ln \left(\frac{\mu^2}{s} \right).$$

Leading coefficients (complex color flow)

Quark-(anti-)quark scattering: $qq \rightarrow qq$ and $q\bar{q} \rightarrow q\bar{q}$

$$c_3 = 2C_F,$$

$$c_2 = -C_F \ln \left(\frac{\mu_\gamma^2}{s} \right) - C_F \ln \left(\frac{\mu_p^2}{s} \right) - \frac{11}{2} C_F$$

$$c_1^\mu = -C_F \left[\ln \left(\frac{p_T^2}{s} \right) + \frac{3}{2} \right] \ln \left(\frac{\mu_p^2}{s} \right) + \frac{\beta_0}{2} \ln \left(\frac{\mu^2}{s} \right)$$

Leading coefficients (complex color flow)

Quark-(anti-)quark scattering: $qq \rightarrow qq$ and $q\bar{q} \rightarrow q\bar{q}$

$$c_3 = 2C_F,$$

$$c_2 = -C_F \ln \left(\frac{\mu_\gamma^2}{s} \right) - C_F \ln \left(\frac{\mu_p^2}{s} \right) - \frac{11}{2} C_F$$

$$c_1^\mu = -C_F \left[\ln \left(\frac{p_T^2}{s} \right) + \frac{3}{2} \right] \ln \left(\frac{\mu_p^2}{s} \right) + \frac{\beta_0}{2} \ln \left(\frac{\mu^2}{s} \right)$$


Similarly for $q\bar{q} \leftrightarrow gg$, $qg \rightarrow qg$, and $gg \rightarrow gg$.

NNLO master formula (simple color flow)

$$\begin{aligned}
 d\sigma_{ab} = & d\sigma_{ab}^B \frac{\alpha_s^2(\mu)}{\pi^2} \left\{ \frac{1}{2} c_3^2 D_3(z) + \left[\frac{3}{2} c_3 c_2 - \frac{\beta_0}{4} c_3 + \sum_j C_{f_j} \frac{\beta_0}{8} \right] D_2(z) \right. \\
 & + \left[c_3 c_1 + c_2^2 - \zeta_2 c_3^2 \frac{\beta_0}{2} T_2 + \frac{\beta_0}{4} c_3 \ln \left(\frac{\mu^2}{s} \right) + \dots \right] D_1(z) \\
 & + \left[c_2 c_1 - \zeta_2 c_2 c_3 + \zeta_3 c_3^2 - \frac{\beta_0}{2} T_1 + \frac{\beta_0}{4} c_2 \ln \left(\frac{\mu^2}{s} \right) + \dots \right] D_0(z) \\
 & \left. + \left[\frac{1}{2} c_1^2 - \frac{\zeta_2}{2} c_2^2 + \frac{1}{4} \zeta_2^2 c_3^2 + \zeta_3 c_3 c_2 + \dots + R \right] \delta(1-z) \right\}
 \end{aligned}$$

Inclusive jet hadroproduction

N. Kidonakis, J. Owens, Phys. Rev. D 63 (2001) 054019 (Fig. 2)



Jet production in DIS

T. Biekötter, M. Klasen, G. Kramer, Phys. Rev. D 92 (2015) 074037

Experimental conditions:

- HERA-II (2003-2007), $\sqrt{S} = 319$ GeV, $\mathcal{L} = 351$ pb $^{-1}$
- $150 \text{ GeV}^2 < Q^2 < 15000 \text{ GeV}^2$, $0.2 < y < 0.7$
- $p_T^{\text{jet}} > 7$ GeV, $-1.0 < \eta^{\text{jet}} < 2.5$, k_T -algorithm with $R = 1$

Jet production in DIS

T. Biekötter, M. Klasen, G. Kramer, Phys. Rev. D 92 (2015) 074037

Experimental conditions:


- HERA-II (2003-2007), $\sqrt{S} = 319$ GeV, $\mathcal{L} = 351$ pb $^{-1}$
- $150 \text{ GeV}^2 < Q^2 < 15000 \text{ GeV}^2$, $0.2 < y < 0.7$
- $p_T^{\text{jet}} > 7$ GeV, $-1.0 < \eta^{\text{jet}} < 2.5$, k_T -algorithm with $R = 1$

Theoretical input:

- Central scales: $\mu^2 = (Q^2 + p_T^2)/2$, $\mu_p^2 = Q^2$
- Proton PDFs: MSTW2008, $n_f = 5$, $\alpha_s(M_Z) = 0.110\dots 0.130$
- Hadronization corrections modeled with PYTHIA


Inclusive jet production in DIS

T. Biekötter, M. Klasen, G. Kramer, Phys. Rev. D 92 (2015) 074037



Dijet production in DIS

H1 Coll., DESY 16-200, to be subm. to EPJC



H1 Dijets

- H1 HERA-II
- Systematic uncertainty
- NLO \otimes hadr. corr.
- aNNLO \otimes hadr. corr.
- NNLO \otimes hadr. corr.

Jet photoproduction

M. Klasen, G. Kramer, M. Michael, Phys. Rev. D 89 (2014) 074032

Experimental conditions:

- HERA-II (2005-2007), $\sqrt{S} = 319$ GeV, $\mathcal{L} = 300$ pb $^{-1}$
- $Q^2 < 1$ GeV 2 , 142 GeV $< W < 293$ GeV
- $p_T^{\text{jet}} > 17$ GeV, $-1.0 < \eta^{\text{jet}} < 2.5$, k_T -algorithm with $R = 1$

Jet photoproduction

M. Klasen, G. Kramer, M. Michael, Phys. Rev. D 89 (2014) 074032

Experimental conditions:


- HERA-II (2005-2007), $\sqrt{S} = 319$ GeV, $\mathcal{L} = 300$ pb $^{-1}$
- $Q^2 < 1$ GeV 2 , 142 GeV $< W < 293$ GeV
- $p_T^{\text{jet}} > 17$ GeV, $-1.0 < \eta^{\text{jet}} < 2.5$, k_T -algorithm with $R = 1$

Theoretical input:

- Central scales: $\mu = \mu_p = \mu_\gamma = p_T$
- Proton PDFs: CT10, $n_f = 5$, $\alpha_s(M_Z) = 0.112\dots 0.124$
- Photon PDFs: GRV-HO, transformed from DIS $_\gamma$ to $\overline{\text{MS}}$
- Hadronization corrections modeled with PYTHIA


Inclusive jet photoproduction

M. Klasen, G. Kramer, M. Michael, Phys. Rev. D 89 (2014) 074032



Inclusive jet photoproduction

M. Klasen, G. Kramer, M. Michael, Phys. Rev. D 89 (2014) 074032



Determination of α_s

M. Klasen, G. Kramer, M. Michael, Phys. Rev. D 89 (2014) 074032

Determination at NLO:

$$\alpha_s(M_Z) = 0.121^{+0.002}_{-0.002}(\text{exp.})^{+0.005}_{-0.003}(\text{th.})$$

Determination of α_s

M. Klasen, G. Kramer, M. Michael, Phys. Rev. D 89 (2014) 074032

Determination at NLO:

$$\alpha_s(M_Z) = 0.121^{+0.002}_{-0.002}(\text{exp.})^{+0.005}_{-0.003}(\text{th.})$$

Determination at aNNLO:

$$\alpha_s(M_Z) = 0.120^{+0.002}_{-0.002}(\text{exp.})^{+0.003}_{-0.003}(\text{th.})$$

Nuclear PDFs from jets in DIS at EIC

EIC White Paper, 1212.1701 [nucl-ex]

eRHIC conditions:

- $E_e = 16 \dots 21$ GeV and $E_A = 100$ GeV $\rightarrow \sqrt{s} = 80 \dots 90$ GeV
- Integrated luminosity: $\mathcal{L} = 10 \dots 3$ fb $^{-1}$

Nuclear PDFs from jets in DIS at EIC

EIC White Paper, 1212.1701 [nucl-ex]

eRHIC conditions:

- $E_e = 16 \dots 21$ GeV and $E_A = 100$ GeV $\rightarrow \sqrt{s} = 80 \dots 90$ GeV
- Integrated luminosity: $\mathcal{L} = 10 \dots 3 \text{ fb}^{-1}$

MEIC conditions:

- $E_e = 12$ GeV and $E_A = 40$ GeV $\rightarrow \sqrt{s} = 45$ GeV
- Integrated luminosity: $\mathcal{L} = 100 \text{ fb}^{-1}$

Nuclear PDFs from jets in DIS at EIC

EIC White Paper, 1212.1701 [nucl-ex]

eRHIC conditions:

- $E_e = 16 \dots 21$ GeV and $E_A = 100$ GeV $\rightarrow \sqrt{s} = 80 \dots 90$ GeV
- Integrated luminosity: $\mathcal{L} = 10 \dots 3 \text{ fb}^{-1}$

MEIC conditions:

- $E_e = 12$ GeV and $E_A = 40$ GeV $\rightarrow \sqrt{s} = 45$ GeV
- Integrated luminosity: $\mathcal{L} = 100 \text{ fb}^{-1}$

Detector specifications:

- Electron or JB method: $Q^2 > 1 \text{ GeV}^2$ and $0.01 \leq y \leq 0.95$
- Electromagn. (hadr.) calorimeter: $-4(-1) < \eta^{\text{jet}} < 4$
- Jet reconstruction in the Breit frame with $p_T^{\text{jet}} > 4 \text{ GeV}$

Nuclear PDFs from jets in DIS at EIC

EIC White Paper, 1212.1701 [nucl-ex]

eRHIC conditions:

- $E_e = 16 \dots 21$ GeV and $E_A = 100$ GeV $\rightarrow \sqrt{s} = 80 \dots 90$ GeV
- Integrated luminosity: $\mathcal{L} = 10 \dots 3$ fb $^{-1}$

MEIC conditions:

- $E_e = 12$ GeV and $E_A = 40$ GeV $\rightarrow \sqrt{s} = 45$ GeV
- Integrated luminosity: $\mathcal{L} = 100$ fb $^{-1}$

Detector specifications:


- Electron or JB method: $Q^2 > 1$ GeV 2 and $0.01 \leq y \leq 0.95$
- Electromagn. (hadr.) calorimeter: $-4(-1) < \eta^{\text{jet}} < 4$
- Jet reconstruction in the Breit frame with $p_T^{\text{jet}} > 4$ GeV

Theoretical input:

- Central scales: $\mu^2 = (Q^2 + p_T^2)/2$, $\mu_p^2 = Q^2$
- Nuclear PDFs: nCTEQ15(-np) with 32 error PDFs


Kinematic acceptance in DIS, DY and at two EICs

EIC White Paper, 1212.1701 [nucl-ex]




Current information from F_2^A/F_2^D

K. Kovarik et al., Phys. Rev. D 93 (2016) 085037







Additional information from inclusive pion data

K. Kovarik et al., Phys. Rev. D 93 (2016) 085037







Inclusive jet production at different EICs

MK, in preparation







Inclusive jet production on different nuclei

MK, in preparation







Sensitivity to nPDFs estimated with nCTEQ15

MK, in preparation



Sensitivity to nPDFs estimated with nCTEQ15-np

MK, in preparation



Conclusion

Theoretical approach:

- Approximate NNLO from threshold resummation
More reliable at higher Q^2 or E_T

Conclusion

Theoretical approach:

- Approximate NNLO from threshold resummation
More reliable at higher Q^2 or E_T

Determination of α_s :

- Final HERA data on 1-/2-/3-jets in DIS and photoproduction
Central α_s value improved, theoretical error reduced → PDG

Conclusion

Theoretical approach:

- Approximate NNLO from threshold resummation
More reliable at higher Q^2 or E_T

Determination of α_s :

- Final HERA data on 1-/2-/3-jets in DIS and photoproduction
Central α_s value improved, theoretical error reduced \rightarrow PDG

Nuclear PDFs from jets at the EIC:

- Kinematic range extends to $Q^2 \leq 10^3 \text{ GeV}^2$ and $x_{\text{Bj.}} \geq 10^{-4}$
Current error shrinks by factor of 5 ... 10, in particular for $f_{g/A}$

Conclusion

Theoretical approach:

- Approximate NNLO from threshold resummation
More reliable at higher Q^2 or E_T

Determination of α_s :

- Final HERA data on 1-/2-/3-jets in DIS and photoproduction
Central α_s value improved, theoretical error reduced \rightarrow PDG

Nuclear PDFs from jets at the EIC:

- Kinematic range extends to $Q^2 \leq 10^3 \text{ GeV}^2$ and $x_{\text{Bj.}} \geq 10^{-4}$
Current error shrinks by factor of 5 ... 10, in particular for $f_{g/A}$

Outlook:

- Improve Kidonakis formalism to account for finite jet mass
D. de Florian, P. Hinderer, A. Mukherjee, F. Ringer, W. Vogelsang,
Phys. Rev. Lett. 112 (2014) 082001
- Full NNLO calculations, e.g. $gg \rightarrow gg$
J. Currie, A. Gehrmann, N. Glover, J. Pires, JHEP 1401 (2014) 110

Jet production in DIS at higher Q^2

D. de Florian, P. Hinderer, A. Mukherjee, F. Ringer, W. Vogelsang, Phys. Rev. Lett. 112 (2014) 082001

